



**HAL**  
open science

# Synthesis of Biologically Relevant 1,2,3-and 1,3,4-Triazoles: From Classical Pathway to Green Chemistry

Lori Gonnet, Michel Baron, Michel Baltas

► **To cite this version:**

Lori Gonnet, Michel Baron, Michel Baltas. Synthesis of Biologically Relevant 1,2,3-and 1,3,4-Triazoles: From Classical Pathway to Green Chemistry. *Molecules*, 2021, 26 (18), pp.5667. 10.3390/molecules26185667 . hal-03360993

**HAL Id: hal-03360993**

<https://imt-mines-albi.hal.science/hal-03360993v1>

Submitted on 1 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Review

# Synthesis of Biologically Relevant 1,2,3- and 1,3,4-Triazoles: From Classical Pathway to Green Chemistry

Lori Gonnet <sup>1,2</sup> , Michel Baron <sup>1</sup>  and Michel Baltas <sup>3,\*</sup>

<sup>1</sup> IMT Mines Albi, UMR CNRS 5302, Centre Rapsodee, Campus Jarlard, Allée des Sciences, Université de Toulouse, CEDEX 09, 81013 Albi, France; lori.gonnet@mcgill.ca (L.G.); michel.baron@mines-albi.fr (M.B.)

<sup>2</sup> Department of Chemistry, McGill University, 801 Sherbrooke St. West, Montreal, QC H3A 0B8, Canada

<sup>3</sup> CNRS, LCC (Laboratoire de Chimie de Coordination), Université de Toulouse, UPS, INPT, Inserm ERL 1289, 205 Route de Narbonne, BP 44099, CEDEX 4, F-31077 Toulouse, France

\* Correspondence: michel.baltas@lcc-toulouse.fr; Tel.: +33-(0)5-61-33-31-15

**Abstract:** Green Chemistry has become in the last two decades an increasing part of research interest. Nonconventional «green» sources for chemical reactions include micro-wave, mechanical mixing, visible light and ultrasound. 1,2,3-triazoles have important applications in pharmaceutical chemistry while their 1,2,4 counterparts are developed to a lesser extent. In the review presented here we will focus on synthesis of 1,2,3 and 1,2,4-triazole systems by means of classical and «green chemistry» conditions involving ultrasound chemistry and mechanochemistry. The focus will be on compounds/scaffolds that possess biological/pharmacophoric properties. Finally, we will also present the formal cycloreversion of 1,2,3-triazole compounds under mechanical forces and its potential use in biological systems.

**Keywords:** 1,2,3-triazoles; 1,2,4-triazoles; ultrasound; medicinal chemistry; green chemistry; mechanochemistry; biological properties



**Citation:** Gonnet, L.; Baron, M.; Baltas, M. Synthesis of Biologically Relevant 1,2,3- and 1,3,4-Triazoles: From Classical Pathway to Green Chemistry. *Molecules* **2021**, *26*, 5667. <https://doi.org/10.3390/molecules26185667>

Academic Editor: José Carlos Menéndez

Received: 5 August 2021

Accepted: 14 September 2021

Published: 18 September 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

One of the main goals in the area of organic synthesis oriented towards biologically active compounds is the research and development of efficient environmentally safe methods. In fact, since the 2000s many regulations for the chemical and pharmaceutical industries have appeared, especially in terms of efficiency, waste management and energy input. All these issues are now addressed and termed «Green Chemistry», a multifaceted field dealing with what we call the twelve principles of P.T. Anastas and J.C. Warner [1]. Most important of them are: atom economy, preventing the use of solvents volatile and/or toxic, minimize chemical waste and minimize energy [2]. Organic reactions and processes are classically conducted in solutions (mostly organic) under reflux or thermal energy to be balanced at the end of the transformation. We focus on the Green Chemistry synthetic aspects, and focus on chemical reactions by using alternative energy sources that appeared and developed since the last two decades; namely, the processes: photochemistry through light excitation, microwave, sonochemistry irradiation, and mechanochemistry [3].

In this article, in order to give an emblematic example of the evolution of synthesis strategies towards ever greener processes, in particular in the pharmaceutical field, we will focus not only on recent classical synthesis of 1,2,3 and 1,2,4-triazoles, but also on sonochemistry and mechanochemical synthesis of these systems in relation to their biological activities. First, we will initially focus on these two alternative energy sources, i.e., sonochemistry/ultrasonic irradiation, and mechanochemistry. Mechanical effects caused by sound irradiation—called sonochemistry—can be applied to liquids. It can induce formation and growth of acoustic cavitations resulting in implosive bubble collapse [4–6]. This leads to intense compressional heating and extremely high pressures in

the resultant so called hot spots (5000 K, 1000 atmospheres) while heating and cooling rates are exceeding  $10^{10} \text{ K s}^{-1}$  (Figure 1) [7,8]. The sonic spectrum ranges from high power to low power ultrasound (20 KHz to 10 MHz). The range from 20 KHz to 1 MHz is used in sonochemistry. As indicated before, ultrasound irradiation can induce formation and growth of acoustic cavitations resulting in implosive bubble collapse that due to their physical properties can substantially improve chemical reactions (catalytic or not) in terms of speed (in some reactions a million fold reactivity increase was observed), selectivity and yield. Ultrasound reactions are not adequate for reactions between solids or solid-gas systems [9]. Ultrasound in organic synthesis has been studied considerably in the past two decades. Especially, various named organic transformations effected through ultrasound irradiation were developed. Among the most important [10] we can point to the coupling reactions, i.e., Heck, Suzuki, Sonogashira, Ullmann, ultrasound-assisted phase transfer catalysis, some named reactions like Reformatsky, Michael, Baylis-Hillmann, but also oxidation/reduction reactions, halogenations. Finally, ultrasound synthesis of ionic liquids and heterocyclic, especially nitrogen contained compounds [11], has gained much success and development. Mechanical energy can also induce chemical transformations [10]. According to IUPAC, a mechanochemical reaction is a “Chemical reaction that is induced by the direct absorption of mechanical energy” [12,13]. Wilhelm Ostwald (Nobel Prize in 1909), was the first who mentioned the term “Mechanochemistry” and defined it as a “branch of chemistry which is concerned with chemical and physico-chemical changes of substances of all states of aggregation due to the influence of mechanical energy”. It is important to mention the pioneering work of Boldyrev et al., on the mechanisms and kinetics in comminuting devices [14–16], serving as a basis for many mechanochemical works. How the absorption of mechanical energy induces chemical transformations in terms of mechanistic understanding is still under investigation and not fully elucidated. Various models were proposed based on solid chemistry knowledge, like “hot spot” and “magma-plasma model” [17–19]. Other also well-known models (spherical, kinetic and impulse . . . ) were equally proposed [20,21]. Many efforts were developed recently towards a mechanistic level understanding of mechanochemical processes [22]. One of the major trends in progress is to research possible links between the mechanical effect and the action of the forces generated at the molecular level [23–27]. In parallel with these recent advances, the topic is still subject to research from the experimental and theoretical points of view [28]. In terms of the experimental view, the traditional grinding by using a mortar and a pestle has been replaced by more sophisticated ball-milling or mechano-milling techniques that are generally conducted in vibration mills or planetary mills at frequencies of 5–60 Hz. The reactions are generally carried out in vessels or jars of different kinds of materials (stainless steel, tungsten carbide, zirconia, agate, etc.). In recent years a deviation from the pure solid status of reactants, named the liquid assisted grinding, gained considerable interest because it offers opportunities to mechanochemistry to reach viable results in comparison to solution synthesis [29,30]. These studies are greatly facilitated by a possible continuous monitoring of mechanochemical reactions [31,32]. One drawback for mechanochemistry is the fact that up to now difficulties exist in practically controlling the air and moisture sensitive reagents. However, Kubota et al. have shown recently [33] that mechanochemistry allows carrying out the syntheses of organometallics sensitive to humidity in air.


In the two last decades, this green chemistry approach has been developed considerably in areas related to inorganic compounds and metal complexes synthesis and related mechanistic aspects [34–36] while less interest was focused on organic mechanochemistry, even after the pioneering work reported by Toda in the 1980s [37] and Kaupp [38]. This is actually changing since the last decade’s focus was essentially on the green chemistry and green processes approach [39]. In recent times, mechanochemical synthetic approaches for creating carbon-carbon, carbon-heteroatom, metal-ligand coordination bonds etc. became important issues and gained considerable attention in the literature [40,41], and many applications were carried out in the field of organic mechanochemistry [42–52]. Among



medicinal chemists (Figure 1). A very recent review treats on those methodologies and on the medicinal attributes of 1,2,3-triazoles [56].


We report here notable current examples (year 2018) of classical synthesis of biologically active compounds bearing this frame, but also recent literature from 2014 concerning synthesis of 1,2,3-triazoles by chemical transformations using alternative energy sources (ultrasonic irradiation).

Alexandre et al. reported in 2018 [57] that compounds based on 4-amino-1,2,3-triazole core as potent inhibitors of indoleamine 2,3-dioxygenase (IDO1) are important targets of immuno-oncology research. The authors screened on a recombinant human IDO1 a library of 350,000 compounds and were able to identify a series bearing the 4-amino-1,2,3-triazole core. Upon chemistry optimisation they obtained compound *N*-(4-chlorophenyl)-2*H*-1,2,3-triazol-4-amine with a remarkable potency ( $IC_{50}$  of 0.023  $\mu$ M) substantially more potent than any other IDO1 inhibitor. Synthesis of this compound differs from all other methods. The synthesis involves diazotization of 4-chloroaniline **1** by sodium nitrite followed by reaction with 2-aminoacetonitrile hydrochloride **2** in order to afford 2-(2-(4-chlorophenyl)iminohydrazino) acetonitrile **3** which upon heating under reflux in ethanol afforded the desired compound **4** (Scheme 1).



**Scheme 1.** Synthesis of 2-(2-(4-chlorophenyl)iminohydrazino) acetonitrile **4**.

Wu et al. [58] reported the design and synthesis of tacrine 1,2,3-triazole derivatives as potent cholinesterase inhibitors. Tacrine **5**, the first drug approved by the FDA that binds at the catalytic active site (CAS) region a potent non selective inhibitor of both bAChE and hBChE was hybridized through various types of linkers bearing the 1,2,3-triazole frame with the chloroquinoline scaffold. Starting from indoline-2,3-dione **6**, the authors obtained in a four step procedure the key tetrahydroacridine intermediates **7** bearing various terminal alkynes. After introduction of an azide functionality in the chloroquinoline scaffold **8** the partners were coupled via CuAAC reaction giving compounds **9** (Scheme 2).




**Scheme 2.** Synthesis of tacrine and quinoline derivatives **9**.

Among all compounds synthesized, compound **9** (R = H, linker = piperazine) exhibited a potent inhibition against AChE and BChE with  $IC_{50}$  values of 4.89 and 3.61  $\mu$ M,


respectively. The authors point out that although this compound is less potent than tacrine, it has a unique binding mode at both CAS and also to the peripheral anionic site (PAS), as well as less toxicity. They concluded by considering it as a lead compound that could be the basis for the development of more active dual inhibitors of AChE and BChE (Table 1).

**Table 1.** Inhibition of AChE (*Electrophorus electricus*) and horse serum BChE by Tacrine 5 and a derivative 9 (with piperazine as linker and R = H).



Compounds	Inhibition (%) at 100 $\mu\text{M}$		$\text{IC}_{50}$ ( $\mu\text{M}$ )	
	AChE	BChE	AChE	BChE
9 (R = H, linker = piperazine)	78.69	91.80	4.89	3.61
Tacrine 5	86.22	99.63	0.316	0.066

Ashok et al. [59] reported the synthesis of a novel prototype that possessed a chromene and a 1,2,3-triazole pharmacophore frame with activities against *M. tuberculosis*. The strategy adopted by the authors for their synthesis started from substituted acetophenone 10 which upon Kabbe condensation and reduction of the carbonyl group afforded spirochromanols 11. Deprotection of 11 and dehydration provided the corresponding spirochromene 12. 1,2,3-triazole-fused spirochromene derivatives 13 were then obtained through a Huisgen cycloaddition in the presence of pyrrolidine as catalyst (Scheme 3).



**Scheme 3.** Synthesis of 1,2,3-triazole-fused spirochromenes 13.


Among the compounds tested against *M. tuberculosis* H37Rv strain, 5 compounds presented strong MIC activities (between 4 and 9  $\mu\text{M}$ ). Their cytotoxicity against RAW 264.7 cells was determined and indicated at least one log difference in comparison to their MIC values. These findings indicated that 1,2,3-triazole-fused spirochromene derivatives can have biological significance for further development (Table 2).

**Table 2.** Anti-tubercular and toxicity evaluation of compounds **13** against *M. tuberculosis* H37Rv.

Compounds <b>13</b>			MIC ( $\mu\text{g/mL}$ )	MIC ( $\mu\text{M}$ )	Cytotoxicity in % Inhibition at 50 $\mu\text{g/mL}$
R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>			
Me	H	C <sub>6</sub> H <sub>5</sub> -	1.56	4.74	30.23
Me	H	4-OMe-C <sub>6</sub> H <sub>5</sub> -	1.56	4.34	33.14
Cl	H	Benzyl	1.56	4.11	29.36
Cl	Me	C <sub>6</sub> H <sub>5</sub> -	3.125	8.60	24.90
Cl	Me	4-Cl-C <sub>6</sub> H <sub>5</sub> -	3.125	7.87	24.76

MIC: minimum inhibitory concentration (the lowest concentration that inhibited the bacterial growth).

López-Rojas et al. [60] reported the synthesis of 4-substituted 1,2,3-triazole coumarin-derivatives and evaluated their antimicrobial activity. The strategy adopted by the authors was the synthesis of acetylenic O- **15** or N-propargylated **17** coumarins starting from 4-hydroxy **14** and 4-bromo **16** coumarin respectively. Copper(I) catalyzed Huisgen 1,3-dipolar cycloaddition reaction with synthesized (or commercially available) alkyl or aryl azides and afforded the desired compounds **18** (Scheme 4).

**Scheme 4.** Synthesis of 4-substituted 1,2,3-triazole-coumarin derivatives **18**.

The authors thus created a focused library of 26 compounds with two isosteric series (hydroxy/amino) and with different substituents at the triazole moiety. Based on their MIC values against selected microorganisms, 5 out of 26 compounds showed significant antibacterial activity towards *Enterococcus faecalis* (MIC = 12.5–50  $\mu\text{g/mL}$ ) while low cytotoxicity was observed against human erythrocytes (Table 3).

**Table 3.** Antimicrobial activity of compounds **18** against *Enterococcus faecalis* (MIC  $\mu\text{g/mL}$ ) of the synthesized compounds.


Compounds <b>18</b>		<i>Enterococcus faecalis</i>
X	R	
O	C <sub>6</sub> H <sub>5</sub> -	50
O	2-OMe-C <sub>6</sub> H <sub>4</sub> -	12.5
O	4-F-C <sub>6</sub> H <sub>4</sub> -	50
NH	3-NO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> -	50
NH	C <sub>11</sub> H <sub>23</sub>	50

### 2.1. Ultrasound Assisted Syntheses of 1,2,3-Triazoles

We will refer herein to some relevant publications from 2014 up to now.

In 2014, Mady et al. [61] reported the ultrasound assisted synthesis of diaryl sulfones bearing 1,2,3-triazole moieties as potential antioxidant and antimicrobial agents. Synthesis of disubstituted triazoles is depicted below (Scheme 5). The authors explored three routes: a stepwise approach that allowed a click coupling of two different azides and a second and third one where both alkynes were introduced then allowed to click from the same azide. Ultrasound (US) Barbier type mediated propargylation occurred readily to construct the common intermediate **19**. This can undergo CuAAC cycloaddition reaction with different

azides under ultrasound conditions affording a first 1,2,3-triazole containing compound **20**. The hydroxy group can be further propargylated and coupled with other azides (or the same), affording final compounds **23**.



**Scheme 5.** Synthesis of mono and bis-1,2,3-triazoles, respectively **20** and **23**.


The second route introduces first a second alkyne group via propargylation of the hydroxy group of key sulfone **21** and then CuAAC cycloaddition reaction with the corresponding azide. All syntheses were operated under ultrasound conditions in a very efficient manner. The authors also synthesized bis-triazoles via the one-pot click reaction (third route).

Biological and antioxidant activities of all compounds were also reported. Many of them were found to be most potent antifungal agents with MIC values around 25  $\mu\text{g}/\text{mL}$  (Table 4). Moreover, compound **24** (Scheme 6) showed an excellent antioxidant activity ( $\text{IC}_{50} = 20 \mu\text{g}/\text{mL}$ ) using a DPPH free radical scavenging assay.

**Table 4.** Minimum inhibitory concentration ( $\mu\text{g}/\text{mL}$ ) against *A. niger*.


Compounds	R	<i>A. niger</i> MIC ( $\mu\text{g}/\text{mL}$ )
<b>20</b>	$\text{C}_6\text{H}_5-$	25
<b>23</b>	$\text{C}_6\text{H}_5-$	25
<b>23</b>	$(4-\text{SO}_2\text{Ph})\text{C}_6\text{H}_4\text{PhCO}-$	25
<b>23</b>	$(4-\text{F})\text{C}_6\text{H}_4\text{PhCO}-$	25
<b>23</b>	$(4-\text{Br})\text{C}_6\text{H}_4\text{PhCO}-$	25





**Scheme 6.** Synthesis of **24** and anti-oxidant activity measured as  $IC_{50}$  by the DPPH procedure.

Nallapati et al. reported in 2015 [62] synthesis of 1,2,3-triazoles derived from olanzapine. Olanzapine (Zyprexa), a member of the thienobenzodiazepine class, is a confirmed marketed drug used for the treatment of schizophrenia and bipolar disorder. The authors describe modifications of olanzapine and explore their activities. One of the target molecules chosen by the authors being olanzapine decorated with 1,2,3-triazole moieties. In that respect alkyne **26** was first prepared through classical coupling in the presence of NaH of propargyl bromide with the drug olanzapine **25** in THF. The thus prepared alkyne reacted with aryl or alkyl azides at room temperature under ultrasound irradiation and in the presence of diisopropylethylamine affording the triazolo derivatives **27** in fairly good yields (Scheme 7).



**Scheme 7.** Synthesis of new olanzapine based 1,2,3-triazole derivatives **27** via CuAAC method.


The authors reported in vitro activities of these compounds against phosphodiesterase 4B protein (PDE4B), a gene family that plays a role in the treatment of schizophrenia. Three of the compounds tested were identified as selective inhibitors of PDE4B ( $IC_{50}$  5 to 6  $\mu$ M) (Table 5).

**Table 5.** Inhibition of PDE4B at 10  $\mu$ M by compounds **27**.

Compounds <b>27</b>	Average % Inhibition against PDE4B
R	
-CH <sub>3</sub> Ph	72.82
-CH <sub>2</sub> C <sub>6</sub> H <sub>3</sub> (Cl- <i>o</i> )(CF <sub>3</sub> - <i>p</i> )	75.37
-C <sub>6</sub> H <sub>4</sub> F- <i>p</i>	74.83

PDE4B: phosphodiesterase 4B protein.

N. Rezki reported in 2015 [63] synthesis under conventional methods and ultrasound conditions of 1,4 disubstituted 1,2,3-triazoles tethering bioactive benzothiazole nucleus and their antibacterial evaluation. Synthesis (Scheme 8) started from 2-aminobenzothiazole derivatives **28** which were acylated upon reaction with bromoacetyl bromide. Then, azidation in the presence of sodium azide afforded the corresponding azidobenzothiazoles **29**. All reactions were performed under classical and ultrasound conditions with better yields in the latter case. Huisgen copper(I) catalysed 1,3-dipolar cycloaddition with appropriate terminal alkynes in the presence of sodium ascorbate in *t*BuOH/H<sub>2</sub>O, and was carried out under heat or use of ultrasound at room temperature affording compounds **30**. Again, ultrasound conditions revealed to be more favorable.



**Scheme 8.** Synthesis of 1,2,3-triazole **30** under ultrasound (US) and conventional methods (CM).

All compounds were tested against three gram positive and three gram negative bacteria and two fungal strains. Some of them, revealed promising activities in the range of 4–8  $\mu\text{g/mL}$  (Table 6).

**Table 6.** Antimicrobial activity of compounds **30** expressed as MIC ( $\mu\text{g/mL}$ ).

Compounds <b>30</b>		Gram-Positive Organisms			Gram-Negative Organisms			Fungi	
R	R <sub>1</sub>	Sp	Bs	Sa	Pa	Ec	Kp	Af	Ca
-SO <sub>2</sub> Me	-	8	8	4	4	8	8	8	8
-SO <sub>2</sub> Me	CH(OH)(Ph)	4	4	8	4	4	8	4	4
-SO <sub>2</sub> Me	-C <sub>2</sub> H <sub>4</sub> OH	4	4	8	4	4	8	4	4

Sp: Streptococcus pneumonia; Bs: Bacillus subtilis; Sa: Staphylococcus aureus; Pa: Pseudomonas aeruginosa; Ec: Escherichia coli; Kp: Klebsiella pneumonia; Af: Aspergillus fumigatus; Ca: Candida albicans.


N. Rezki and M.R. Aouad reported in 2017 [64] synthesis of hybrid compounds bearing fluorinated 1,2,4-triazole, 1*H*-1,2,3-triazole and also a benzothiazole functionality. Construction of the 1,2,4-triazole substituted frame started from reaction of 2-fluorobenzoyl chloride **31** with hydrazine hydrate.

Subsequent treatment was administered with diverse alkyl/aryl isothiocyanates, which upon basic reaction conditions underwent an oxidative ring closure affording the thione derivatives **32**. The latter reacted with propargyl bromide in the presence of triethylamine under ultrasound conditions, furnishing the thiopropargylated 1,2,4-triazole precursors **33** required for the click reaction. On the other hand, acylation of the appropriate 2-aminobenzothiazoles **34** followed by the azidolysis reaction allowed obtention of the azidoacetamide derivative **35**. The Huisgen cycloaddition reaction was then performed between the two coupling reagents in the presence of CuSO<sub>4</sub> and Na-ascorbate as catalysts in DMSO-H<sub>2</sub>O. The ultrasound conditions were less time consuming and much more efficient with almost quantitative yields (Scheme 9).


Almost all compounds showed activities with MIC values in the range 6.45–33.2  $\mu\text{mol/L}$  against *S. pneumoniae*. In addition, compound **37** (Figure 2) showed the strongest antifungal activities among all compounds with MIC values of 6.45  $\mu\text{mol/L}$  against *A. fumigatus* and *C. albicans*.

## 2.2. Mechanochemical Syntheses of 1,2,3-Triazoles


Praveen et al. reported in 2017 [65] the synthesis of new hybrid pharmacophores under ball milling conditions through two well established named reactions, namely a Baylis-Hillman [66] and a Huisgen's click chemistry [67]. The authors aimed to prepare potential medicinal targets bearing a 3-substituted-3-hydroxy-2-oxindole frame present in many natural products and medicinal agents [68–70] and a 1,2,3-triazole scaffold. The authors successfully combined a Baylis-Hillman and a click reaction by using DABCO as a base and copper oxide nanoparticles as catalysts. By milling together a mixture of *N*-propargyl isatin, *N*-methylmaleimide, benzyl azide, DABCO in the presence of CuONP catalyst (5%) they were able to find optimal conditions of achieving the synthesis of the target compound **38a** in 96% of yield (Scheme 10).



**Scheme 9.** Synthesis of fluorinated-1,2,4-triazole **36**.



**Figure 2.** 2-(4-(((3-(2-fluorophenyl)-4-phenyl-4,5-dihydro-1H-1,2,4-triazol-5-yl)thio)methyl)-1H-1,2,3-triazol-1-yl)-N-(5-(methylsulfonyl)-2,3-dihydro-1H-inden-2-yl)acetamide **37**.




**Scheme 10.** Mechanochemical synthesis of oxindole-triazole hybrid pharmacophores **38**.

They then applied the conditions found for creating a small focused library (Table 7) as their methodology and accommodated a large variety of substituted starting compounds. In addition, the authors proved the recyclability of the catalyst and its total recovery. Biological studies along with molecular docking demonstrated the rational efficiency of the compounds as antibacterial and antifungal. The best activities were found for compound **38m** (Figure 3), which was most active against *S. aureus* (with a MIC value of  $16 \mu\text{g mL}^{-1}$ ), for compounds **38a,d,i,l** active against *E. coli* and for compounds **38e,h,k,m,p** active against *C. albicans*.


**Table 7.** Compounds **38**: Structures and mechanochemical yield.

Compounds <b>38</b>				Yield (%)
	R	R <sub>1</sub>	R <sub>2</sub>	
<b>38a</b>	Me	H	benzyl	96
<b>38b</b>	Me	H	p-methylbenzyl	98
<b>38c</b>	Me	H	p-methoxybenzyl	99
<b>38d</b>	Me	H	hexyl	80
<b>38e</b>	Me	H	phenyl	70
<b>38f</b>	Me	H	ethoxycarbonylmethyl	90
<b>38g</b>	Me	H	p-nitrobenzyl	93
<b>38h</b>	Me	5-Cl	benzyl	99
<b>38i</b>	Me	5-Cl	p-methylbenzyl	90
<b>38j</b>	Me	5-Cl	p-nitrobenzyl	90
<b>38k</b>	Me	5-Cl	p-methoxybenzyl	98
<b>38l</b>	Me	5-Cl	phenyl	71
<b>38m</b>	phenyl	H	benzyl	90
<b>38n</b>	benzyl	H	benzyl	93
<b>38o</b>	Me	5-Br	benzyl	96
<b>38p</b>	Me	5-Me	benzyl	88



**Figure 3.** Most active compound **38m** of the series.

Sahu et al. published in 2019 [71] the synthesis of quinine-triazole systems with the aim to find new compounds via molecular hybridization [72] that can be addressed to two antiprotozoal targets that are malaria and leishmaniasis [72,73]. The synthetic route (Scheme 11) started with activation through mesylation of the secondary alcohol of quinine generating the compound **39** and subsequent substitution with the azide group via a solution-based methodology [74]. The generated azido dehydroxyquinine **40** was then allowed to react via a copper catalyzed cycloaddition reaction with a variety of alkynes. These reactions were carried out under mechanochemical conditions in ball mill at 300 rpm, affording the triazolyl compounds **41** in 45% to 91% yields (Table 8). Screening results showed that from the 19 synthesized compounds, 5 showed significant antimalarial and antileishmanial activities (Table 8) and four of them did not reveal any in vivo (rodent animal model) toxic manifestation at doses as high as 1000 mg/Kg.




Scheme 11. Synthesis of triazolyl compounds from Quinine.

Table 8. Compounds 41 Structure, yield and activity.

Compounds 41	R	Yield (%)	Antimalarial Assay (IC <sub>50</sub> , (μM))	Antileishmanial Assay (IC <sub>50</sub> , (μM))	Toxicity Profile NOAEL (mg/kg)
a	—C <sub>6</sub> H <sub>13</sub>	91	0.25	8.26	>1000
b	—C <sub>14</sub> H <sub>29</sub>	89	0.25	4.4	>1000
c		56	0.25	1.78	>1000
d		82	0.37	3.95	>1000
e	—CH <sub>3</sub>	78	0.76	4.06	

Finally, S. Sampath et al. reported last year [75] the synthesis of 1,2,3-triazole tethered 3-hydroxy-2-oxindoles under ball milling conditions (Scheme 12) as corrosion inhibitors and antimicrobials. 3-Functionalized oxindoles can be obtained from the valuable heterocyclic scaffolds isatins. Among the different 3-substituted oxindoles, the 3-hydroxy-3-substituted-2-oxindoles are present in many natural products [76]. In addition, they are considered as valuable key intermediates in organic synthesis [77–79], leading to compounds with pronounced pharmaceutical properties [80–82]. The authors synthesized a set of new derivatives of this family by combining an aldol condensation and a click reaction using ball milling conditions. A mixture of N-propargyl isatin, acetophenone, benzyl azide in the presence of DABCO and copper oxide nanoparticles CuONPs (2.5 mol%) was reacted in a ZrO<sub>2</sub> jar material at a speed of 400 rpm, affording the desired products **42** in 87% to 92% yields, except for azide possessing the strong electron withdrawing NO<sub>2</sub> group (80% yield). Among the compounds synthesized, derivative **42b** displayed a remarkable corrosion inhibition potency (for corrosion inhibition in acidic media see references [83,84]), while compound **42a** showed appreciable antifungal (*C. albicans*) and antibacterial (*S. aureus*) effects (Table 9). The authors consider that the biological results are quite encouraging for triggering a detailed structure-activity study and the comprehension of their activity.




**Scheme 12.** Synthesis of 1,2,3-Triazole Tethered 3-Hydroxy-2-oxindoles.

**Table 9.** Compounds 42 Structure, Yield and Activity.

Compounds 42	R	R <sub>1</sub>	R <sub>2</sub>	Yield (%)	Biological Effect
a		H		90	Anti-fungal Anti-bacterial
b		5-Me		92	Main activity (f.i.): Corrosion inhibition

### 3. 1,2,4-Triazole Systems

The 1,2,4-triazole-based biologically active compounds have found enormous applications in medicinal and agricultural sciences. A great number of drugs are extensively used in clinics. Among them, we can point to the antifungal fluconazole **43**, antitumoral letrozole **44**, and the antiviral ribavirin **45**, (Figure 4) [85], while several triazole based compounds play an important role in agriculture ensuring harvest and crops [86]. Their extensive medicinal, agrochemical potential, resulted in an overwhelming effort to develop synthetic methods that include three categories of synthetic objectives: (a) cyclizations to form the triazole ring, (b) transformations of heterocyclic compounds to construct the triazole ring, and (c) substitutions on the 1,2,4-triazole ring.




**Figure 4.** Antifungal fluconazole **43**, antitumoral letrozole **44**, the antiviral ribavirin **45**.

In this review, we discuss only some examples of the recent cyclization reactions with amidrazones and hydrazides. In addition, it is noteworthy to point out that there are no reported methods to synthesize 1,2,4-triazole frames under green chemistry conditions. We report here our first results concerning the mechanochemical organic synthesis of a valuable annulated 1,2,4-triazole scaffold.

### 3.1. From Amidrazones


Amidrazones are the conjugated products of imines and hydrazines; their cyclisation with carbonyl compounds is one of the most important pathways to access 1,2,4-triazole derivatives.

In 2015, Nakka et al. [87] reported an environmentally benign protocol for the synthesis of 3,4,5-trisubstituted 1,2,4-triazoles **48**. The authors performed the coupling/cyclisation reaction by heating in polyethylene glycol and in the presence of ceric ammonium nitrate (catalyst, 5%), *N*-arylamidrazones **46** and aldehydes **47**. The authors demonstrated that this protocol could generate good yields of 3,4,5-tri-substituted 1,2,4-triazoles bearing different functionalities, while in addition, the effective recyclability of the medium could make the process industrially interesting (Scheme 13).




**Scheme 13.** Synthesis of 3,4,5-trisubstituted 1,2,4-triazoles **48**.

Szöcs et al. [87] have already shown that 1,2,4-triazole frames judiciously attached at the 5-position of a 3-*C*-glucopyranosyl scaffold give access to very efficient inhibitors of glycogen phosphorylase. In that respect they become extremely important hits as potential antidiabetic agents (type 2 diabetes). Szöcs et al. reported in 2015 [88] 38 new developments concerning the synthetic approaches for this class of compounds and in particular the oxidative ring closures of *N*<sup>1</sup>-alkylidene carboxamidrazones. When glycosyl cyanides **49** and amidrazones **50** were treated under Raney reductive conditions in the presence of NaHP<sub>2</sub>O<sub>2</sub>, they afforded the two tautomeric forms **51** of *O*-peracylated *N*<sup>1</sup>-(β-*D*-glucopyranosyl-methylidene)-arenecarboxamidrazones. Bromination of **52** by *N*-bromosuccinimide (NBS) led to halogenated **47** type derivatives. The latter can then undergo in pyridine or by NH<sub>4</sub>OAc in AcOH, a ring closing reaction to the desired 3-*C*-glycosyl-5-substituted-1,2,4-triazoles **53** (Scheme 14).



**Scheme 14.** Synthesis of 3-*C*-glycosyl-5-substituted-1,2,4-triazoles **52**.


The authors took advantage of their methodology to create 3,5-diaryl-1,2,4-triazoles **55** starting from *N*<sup>1</sup>-arylidene-arenecarboxamidrazones **54**. Reaction of the latter with NBS/NH<sub>4</sub>OAc in AcOH (whatever the order) afforded triazoles **55**, thus demonstrating the general applicability of the method (Scheme 15).



$Ar_1 = Ph, 2\text{-Pyridyl}$


**Scheme 15.** Syntheses of  $N^1$ -arylidene-arenecarboxamidrazones and their transformation into 3,5-diaryl-1,2,4-triazoles **55**.

Among the different compounds synthesized and tested, it is noteworthy to point out that compound **56** (Figure 5) with an inhibition constant  $K_i$  of  $0.41 \mu M$  against rabbit muscle glycogen phosphorylase could be considered as the starting point for the development of more potent compounds for pharmacological treatment, not only of diabetes but also wherever the regulation of glycogen metabolism plays a significant role (cerebral and cardiac ischemias, and tumor growth).



**Figure 5.** C-glucopyranosyl 1,2,4-triazoles **56**.

In 2018, Aly et al. [89] reported a general method for the synthesis of 1,3,5-trisubstituted 1,2,4-triazoles **59**, **60** from reaction of amidrazones **57** with diethyl azodicarboxylate **58**. The authors performed the coupling/cyclisation reaction between  $N$ -arylamidrazones **57** and diethyl azodicarboxylate **58**. The reaction was conducted under reflux in EtOH and catalyzed by a few drops of triethylamine (Scheme 16), thus allowing to get an easy access to the highly diverse triazoles **59**, **60**. The reaction is based on oxidation of ethanol to acetaldehyde via the Mitsunobu reagent; upon reaction with amidrazones the substituted 3-methyltriazoles **59** could be obtained, while a [2 + 3] cycloaddition reaction between two oxidized forms of amidrazones produced triazoles **60**.




**Scheme 16.** Reaction of the Mitsunobu reagent on amidrazones **57**.

### 3.2. From Hydrazides

Several hydrazides are commercially available and the non-commercial ones are successfully prepared by the reaction of hydrazine with the corresponding ester precursor. Up to now, a lot of works have been done concerning the cyclization of hydrazides or their derivatives. Very recent work in relation to biological activities is presented here.

In 2018, Singh et al. [90] reported the design and synthesis of new bioactive 1,2,4-triazoles as potential antituberculosis and antimicrobial agents. In that respect, they synthesized a series of functionalized 1,2,4-triazole derivatives through the synthetic scheme presented below (Scheme 17).





Scheme 17. Synthesis of 1,2,4-triazole **64** from hydrazides.

Isonicotinic acid hydrazide **61** was transformed to the potassium dithiocarbazinate **62** by reaction with carbon disulfide under basic conditions. Then, treatment with hydrazine hydrate under thermal conditions in water, afforded the 4-amino-1,2,4-triazole-3-thiol **63**. The latter reacted on its 4-amino group to form various Schiff base compounds. Some of the compounds were found to have very potent antitubercular activities, even better than isoniazid and also against clinical isolates (Table 10).

Table 10. Percent reduction in relative light units (RLU) of compounds **64** against *M. tuberculosis* H37RV and clinical isolates, S, H, R and E resistant *M. tuberculosis* at 100 µg/mL.

Compounds <b>64</b>	Percent Reduction in RLU		
	R	<i>M. tuberculosis</i> H <sub>37</sub> Rv	Clinical Isolates, S, H R and E resistant <i>M. tuberculosis</i>
4-F-C <sub>6</sub> H <sub>4</sub> -		80.50	50.72
4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> -		82.01	52.79
CH=CH-C <sub>6</sub> H <sub>5</sub>		83.49	52.47


Synthesized compounds were also tested in vitro against representative bacterial and fungi strains, one compound has very potent activity against *B.subtilis*, while another one is very potent against *A.niger* and *C. albicans* fungi (Table 11).

Table 11. In vitro antibacterial and antifungal activities of compounds **64**.

Compounds <b>64</b>	Minimum Inhibitory Concentration (MIC, µg/mL)				
	R	Gram-Positive Bacteria		Fungi	
		<i>S. aureus</i>	<i>B. subtilis</i>	<i>A. niger</i>	<i>C. albicans</i>
4-F-C <sub>6</sub> H <sub>4</sub> -	12.5	10.2	125	106.3	
-CH=CH-C <sub>6</sub> H <sub>5</sub>	75	81.3	11.7	10.9	

Sonawane et al. reported in 2017 [91] the synthesis of 1,2,4-triazole-3-thione derivatives as antimycobacterial agents. The two routes employed by the authors are outlined in Scheme 18. The acid chloride **66**, prepared by reacting aromatic carboxylic acid **61** with thionyl chloride reacted with thiosemicarbazide, which without isolation and upon thermal heating under aqueous basic conditions, led to the desired compounds **67** (route A). Triazolethiones **69** could not be synthesized by this procedure were obtained by reaction of hydrazide **68** with carbon disulfide, followed by heating in the presence of a 25% ammonia solution (route B).


Two of the compounds synthesized showed high antitubercular activity against the dormant H37Ra strain in vitro and ex vivo; they also showed extremely low cytotoxicity and high solubility indicating the potential of developing these compounds further as novel therapeutics against tuberculosis infection (Table 12).

Scheme 18. Synthesis of 1,2,4-triazole-3-thione **67** and **68**.Table 12. In vitro anti-tubercular and anti-bacterial (gram-positive and gram-negative bacteria) activities of thione derivatives **67**, **69** (IC<sub>50</sub> and MIC values—μg/mL).

Compound	Anti-Tubercular Activity against MTB (μg/mL)		Cytotoxic Activity of Triazole Thiones against Human Cancer Cell Lines (μg/mL)			Aqueous Solubility (μM)
	Dormant Stage	Active Stage	THP-1	A549	PANC-1	
	MIC	MIC	GI <sub>90</sub>	GI <sub>90</sub>	GI <sub>90</sub>	
	0.64	9.05	>100	>100	>100	>1280
	0.46	>30	>100	>100	>100	>3200


MTB H37Ra, *Mycobacterium tuberculosis* H37Ra; Growth Inhibition (GI): GI<sub>90</sub> (concentration which resulted in 90% decrease in cell viability). Expressed in μg/mL. THP-1: acute monocytic leukemia; A549: lung adenocarcinoma; PANC-1: pancreas carcinoma.

Liu et al. [92] reported in 2017 a family of 7-hydroxy-4-phenylchromen-2-linked 1,2,4-triazoles with potent antitumoral activities. The synthetic procedure adopted made use of the coumarin synthesized derivatives **70** that were functionalized by reaction with ethylbromoacetate followed by transformation of the ester group to a hydrazide functionality. The latter, when condensed with dimethylacetal followed by a strong thermal reaction with an amine in the presence of glacial acetic acid afforded the triazole derivatives **73** in good overall yields (Scheme 19).

Scheme 19. Synthesis of 7-hydroxy-4-phenylchromen-2-linked 1,2,4-triazoles **73**.

The new 1,2,3-triazole derivatives showed improved antiproliferative activities. Particularly, compound **74** exhibited potent activity with important IC<sub>50</sub> values against AGS (2.63 ± 0.17), MGC-803 (3.05 ± 0.29) and HCT-116 cell lines (11.57 ± 0.53 μM) (Figure 6). The authors also demonstrated that these compounds had strong activity against the HeLa cell line, with an IC<sub>50</sub> value of 13.62 ± 0.86 μM. All activities were better than those of the non-substituted 7-hydroxy-4-phenyl-2H-chromen-2-one **70** (with R = H) and that of the positive control drug 5-fluorouracil. Moreover, the authors showed that the biolog-

ical activities of the 1,2,4-triazole derivatives were significantly higher than that of the 1,2,3-triazole ones.




**Figure 6.** 7-hydroxy-4-phenyl-2H-chromen-2-one **70** (with R = H) and 7-((4-(4-Chlorophenyl)-4H-1,2,4-triazol-3-yl)methoxy)-4-phenyl-2H-chromen-2-one **74**.

### 3.3. Annulated 1,2,4-Triazole Systems

Among the annulated 1,2,4-triazole systems we will present the triazolophthalazine frame that was developed in our group by conventional and non-conventional means.


Some years ago, De et al. [93] explored the possibility of cinnamic acid derivatives as potential antituberculosis agents. In the course of their studies the authors synthesized 4-alkoxy cinnamoyl derivatives resulting from the coupling of the corresponding acids (or activated ones) with different nucleophiles and among them amines, hydrazines, thiols.

In the course of their first studies, when reacting under peptide coupling conditions (EDC, HCl, HOBT, and trimethylamine), 1-hydrazinophthalazine hydrochloride **75**, and cinnamic acid derivatives for 48 h under reflux in acetonitrile, the authors obtained in good yields the corresponding 3-(4-alkoxystyryl)-[1,2,4]triazolo[3,4-a]phthalazines **77** (65–90%). This was formed through a coupling-intramolecular cyclization-dehydration sequence (Scheme 20).




**Scheme 20.** Synthesis of 3-(4-alkoxystyryl)-[1,2,4]triazolo[3,4-a]phthalazines **77**.

All alkoxyated compounds showed good antitubercular activities. More importantly, triazolophthalazine derivative **73** (Figure 7), bearing a 4-isopentenylxy chain on the phenyl ring, showed excellent antitubercular potency (MIC = 1.4 μM) in addition to a very good cytotoxicity toward HCT116 human cells (IC<sub>50</sub> = 449 μM; 160 μg/mL and selectivity index SI = 320). It is also noteworthy to point out that this compound does not act on the mycolic acid biosynthesis of mycobacteria and up to now its target is unknown.




**Figure 7.** (E)-3-[4-(3-Methylbut-2-enyloxy)styryl]-[1,2,4]triazolo[3,4-a]phthalazine **78**.

In order to build a small, focused library of styryltriazolophthalazines, Veau et al. [94] modified the convergent route to a divergent one, by exploring the possibility of the construction of the phenolic precursor **79** that could then lead to various alkoxyated derivatives of type **77** (Scheme 21).



**Scheme 21.** Convergent (A) and divergent (B) routes for synthesis of compounds 77.


The target precursor **79** was thus obtained by the authors in a two step procedure: 1-hydrazinophthalazine hydrochloride **74** and *p*-hydroxy cinnamic acid **80** when reacted in acetonitrile for 1 h under peptidic coupling conditions but under microwave afforded a 69% yield after a simple filtration–recrystallization sequence of the styrylphenolic triazolophthalazine **79**. Alkylation by various alkylating agents and under standard conditions led to the desired alkoxyated derivatives **81** (alkylating agent,  $\text{K}_2\text{CO}_3$ , KI, DMF,  $60^\circ\text{C}$ , overnight) (Scheme 22).



**Scheme 22.** Synthesis of alkoxyated *trans*-styryltriazolophthalazine derivatives **81**.


In continuation of this work on the triazolophthalazine frame the authors considered that this could be an interesting pharmacophore to explore. In that respect they also synthesized two compounds bearing either an alkyne group or a bromine at the 2 position of the 1,2,4-triazole frame.

Concerning the alkyne compound, after several attempts, Veau et al. [94] considered the best way to obtain it is a two-step sequence. Coupling of trimethylsilyl propiolic acid **82** with 1-hydrazinylphthalazine hydrochloride **74** under mild conditions afforded acyclic precursor **83**. Under microwave irradiation in acetonitrile for 1 h, the precursor was transformed quantitatively to the cyclized silylated intermediate **84**. Upon standard deprotection conditions (i.e.,  $\text{K}_2\text{CO}_3$  in MeOH) compound **84** led to derivative **85** (31% overall yield over three steps) (Scheme 23).




**Scheme 23.** Synthesis of 3-ethynylated derivative **85**.

For the first time, Gonnet et al. [95] reported in 2019 the mechanochemical synthesis of 1,2,4-triazoles starting from hydralazine hydrochloride. By using a planetary ball-mill, and in the presence of pyrogenic S13 silica as the grinding auxiliary, total conversion to intermediate hydrazones **86** was achieved in a few minutes (Scheme 24). Iodobenzene diacetate (IBD) was used for optimal conversion of nonphenolic hydrazones to annulated 1,2,4-triazoles **87**, while SeO<sub>2</sub> was found to be efficient for phenolic compounds (Scheme 25).



**Scheme 24.** One pot two step mechanochemical synthesis of annulated 1,2,4-triazoles **87**.



**Scheme 25.** One pot two step synthesis of phenolic 1,2,3-triazoles **89**.

In addition, for the first time, the one-pot two-step synthesis (Scheme 24) leading to annulated 1,2,4-triazoles was also successfully conducted.


Comparison to the conventional syntheses of hydrazone **86** and triazole **87** clearly showed the green metrics overall efficiency of the mechanochemical synthesis (Table 13).

**Table 13.** Comparison of Green metrics for the synthesis of Hydrazone **86** and Triazole **87**, in solution/by mechanochemistry.

Compound	Time (h)	Yield (%)	E-factor	PMI
<b>86</b>	1/0.5	100/100	14/4	15/5
<b>87</b>	4/0.75	84/97	74/12	75/13

Synthesis of the brominated compound **91** was also explored. First a conventional method was performed by Veau et al. [94], by reacting overnight under reflux in ethanol, 1-hydrazinyl-phthalazine hydrochloride **74** with trimethylortho-formate in the presence of some drops of acetic acid in order to obtain unsubstituted triazolophthalazine **90**. Then, reaction with bromine in the presence of pure acetic acid under reflux afforded the brominated compound **91** in 53% yield.

More recently, Gonnet et al. [95] reported the two-step bromination reaction successfully conducted by mechanochemistry. Reaction of 1-hydrazinylphthalazine hydrochloride **74** in the presence of trimethylorthoformate and some drops of acetic acid reacted in a planetary ball mill for 1 h affording quantitatively triazolo-phthalazine **90**. Reaction in the same planetary ball-mill (PBM) of the triazolophthalazine with sodium bromide, oxone and some silica afforded after 1 h the brominated compound **91** in quantitative yield (Scheme 26).




**Scheme 26.** Syntheses of 3-bromo-[1,2,4]triazolo[3,4-a]phthalazine **91** by classical method (above) and mechanochemical condition (below).

All compounds bearing the 1,2,4-triazole frame were evaluated for various biological properties. The 3-aryl substituted 1,2,4-triazole derivatives **87** and **89** do not present valuable activities against *M.tuberculosis* (MIC around 80  $\mu\text{M}$ ) [96]. The alkyne derivative **85** present a very good activity (MIC 12.9  $\mu\text{M}$ ) while the brominated **91** is much less potent (MIC 40  $\mu\text{M}$ ) [94]. The alkyne derivative **85** did not manifest cytotoxicity toward HCT116 human cells while it is equally active against multidrug-resistant *M. tuberculosis* strains. Considering all results starting from compound **73**, it seems likely that the triazolophthalazine could be an important scaffold in order to obtain new families of compounds with strong antitubercular activity and an alternative mode of action for compared with standard anti *M. tuberculosis* drugs.

In that respect, the authors consider the possibility of developing focused libraries of triazolophthalazine compounds by using the two important precursors that are the alkyne and the brominated derivatives and developing coupling reactions under conventional and/or mechanochemical means. In addition, further work is necessary to tackle the identification of the protein targeted by this class of potent anti *M. tuberculosis* compounds.

#### 4. Mechanochemical Cycloreversion of 1,2,3-Triazoles


Globally, the cycloaddition process is strongly favored thermodynamically ( $\Delta H = -45$  to  $-55$  Kcal/mol) [97]. The 1,2,3-triazole frame is robust and inert under most thermal chemical treatments but also in aqueous or biological environments. In 2011, Brantley et al. [98] reported the possibility of unclicking the click on specific 1,4-substituted 1,2,3-triazoles by mechanical forces. They first hypothesized that mechanical exogenous forces directed to judiciously chosen scaffolds incorporated in a polymer chain can formally disallow pericyclic reactions. The authors incorporated the triazole ring in polymer chains and one of them was judiciously chosen, when mechanical ultrasound forces were applied (ultrasonication in a Suslick cell at  $0^\circ\text{C}$ ) resulted in the cleavage of the triazole ring **92** to its alkyne **93** and azide **94** components (Scheme 27).



**Scheme 27.** Effect of ultrasound on the triazole **92**.

They concluded that the ability to selectively deconstruct triazoles might serve to elaborate mechano-responsive materials for potential controlled bioconjugation applications or force responsive fluorescent tags for biological assays.

This stimulated a puzzling publication (that was retracted since by the editor) [99] aroused strong debate. The same authors, based on sonochemical experiments related to extended Bel theory, discussed and concluded on the lowering of the activation energy barrier for cycloreversion [100] through application of an external force to the triazole ring. In the contrary, purely theoretical work, it was shown that the cycloreversion barrier is as high as 70 Kcal/mol [101]. In addition, the mechanochemically induced retro-click of the 1,2,3-triazole ring vs. bond rupture next to it could not be unambiguously concluded when single molecule force spectroscopy experiments were applied [100]. Stauch and Drew reported in 2017 [102] a theoretical work where by using the JEDI (Judgment of Energy DIstribution) analysis it was concluded that for 1,4 disubstituted triazoles the unclick reaction is impossible, even when  $\text{Cu}^{\text{I}}$  assisted (Scheme 28a). For 1,5-disubstituted triazoles where a parallel alignment of the scissile bond exists, this could be feasible. Nevertheless, the retro click cycloreversion is not selective as it competes with the carbon-nitrogen bond connecting the triazole ring to the linker. During the same year, Krupička et al. [103] also concluded that only in these 1,5-disubstituted 1,2,3-triazole systems are the Gibbs free energy barriers 55 Kcal/mol (unclick reaction) versus 45 Kcal/mol for external C-N bond cleavage. The authors also point out an extremely exciting finding by showing that the calculated Ru-assisted mechanochemical cycloreversion of the 1,5 regioisomer dramatically lowers the activation energy of the rate determining step down to 20 Kcal/mol (the first step), while the decomplexation of the cleaved intermediate readily occurs, leading to the alkyne and azide components (Scheme 28b).



**Scheme 28.** Reactions of (a) 1,4-dimethyl 1,2,3-triazole leading to mononuclear Cu<sup>I</sup>-catalyzed cycloreversion and (b) 1,5-dimethyl 1,2,3-triazole leading to Ru<sup>II</sup>-catalyzed cycloreversions.

In conclusion, the authors point out that the Ru-assisted mechanochemical unclicking of the 1,5 regioisomer could be an extremely selective process. If this is to be experimentally proved it would open the path for very important potential applications.

## 5. Conclusions

Among the nitrogen-contained heterocyclic ring structures one of the most important providing long term advancement in the medical field are triazoles. They became in the last decades the heterocycle of choice in all fields of drug discovery receiving much of the attention and offering new opportunities for medicinal chemists.

We must point out that while 1,2,3-triazole systems are very well documented in terms of classical organic synthesis, their synthetic methodologies under green chemistry approaches based on less energy input requirements are beginning to emerge, but are still focused on ultrasound reactions. It is our feeling that the mechanical approaches will be further developed (mechanochemical synthesis, ultrasound). There is still a lot to be invented and this is a great opportunity for the chemists and medicinal chemists, but also in extenso for the pharmaceutical industry.

The same and in a greater extent is also true for the synthesis of the valuable regioisomeric scaffold of 1,2,4-triazole systems. Except for our contribution in the field, there is no other green chemistry (ultrasound or mechanochemistry) developed for this family of compounds. We believe that here also great opportunities exist for all communities of synthetic, physical, theoretical and medicinal chemists, whether they are in the academia or in the industry.

Finally, concerning the mechanochemical (ultrasound) unclicking of the 1,5 disubstituted 1,2,3-triazole, active experimental research work is needed in order to create an extremely selective process that could also confirm the theoretical work. That could pave the way for important biological and other applications.

**Author Contributions:** L.G.: resources, writing, review and editing. M.B. (Michel Baron): resources, writing, review and editing. M.B. (Michel Baltas): Conceptualization, writing—original draft preparation, review and editing. All authors have read and agreed to the published version of the manuscript.

**Funding:** The authors would like to gratefully acknowledge the Region Occitanie for the thesis grant (L.G.) and the CNRS, University Paul Sabatier, IMT Mines Albi for financial support.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Sample Availability:** Samples of the compounds are not available from the authors.



## References

1. Anastas, P.T.; Warner, J.C. *Green Chemistry, Theory and Practice*; Oxford University Press: Oxford, UK, 2000; ISBN 0198502346/9780198502340.
2. Poliakov, M.; Licence, P. Sustainable Technology: Green Chemistry. Available online: <https://www.nature.com/articles/450810a> (accessed on 16 September 2018).
3. Varma, R.S. Chemical Activation by Mechanochemical Mixing, Microwave and Ultrasonic Irradiation. *Green Chem.* **2008**, *10*, 1129–1130. [[CrossRef](#)]
4. Mason, T.J.; Lorimer, J.P. *Applied Sonochemistry: Uses of Power Ultrasound in Chemistry and Processing*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2002; ISBN 978-3-527-60054-0.
5. Bonrath, W.; Schmidt, R.A.P. Ultrasound in Synthetic Organic Chemistry. *Adv. Org. Synth.* **2005**, *1*, 81–117. [[CrossRef](#)]
6. Li, J.T.; Wang, S.X.; Li, G.F.C.; Li, T.S. Some Applications of Ultrasound Irradiation in Organic Synthesis. Available online: <http://www.eurekaselect.com/79542/article> (accessed on 16 September 2018).
7. Didenko, Y.T.; Suslick, K.S. The Energy Efficiency of Formation of Photons, Radicals and Ions during Single-Bubble Cavitation. *Nature* **2002**, *418*, 394–397. [[CrossRef](#)] [[PubMed](#)]
8. Suslick, K.S.; Flannigan, D.J. Inside a Collapsing Bubble: Sonoluminescence and the Conditions during Cavitation. *Annu. Rev. Phys. Chem.* **2008**, *59*, 659–683. [[CrossRef](#)] [[PubMed](#)]
9. Xu, H.; Zeiger, B.W.; Suslick, K.S. Sonochemical Synthesis of Nanomaterials. *Chem. Soc. Rev.* **2013**, *42*, 2555–2567. [[CrossRef](#)] [[PubMed](#)]
10. Baig, R.B.N.; Varma, R.S. Alternative Energy Input: Mechanochemical, Microwave and Ultrasound-Assisted Organic Synthesis. *Chem. Soc. Rev.* **2012**, *41*, 1559–1584. [[CrossRef](#)]
11. Gaudino, E.; Tagliapietra, S.; Mantegna, S.; Cravotto, G. Mechanochemical and Sonochemical Heterocyclizations. *Chem. Heterocycl. Compd.* **2016**, *52*, 856–865. [[CrossRef](#)]
12. Stauch, T.; Dreuw, A. Advances in Quantum Mechanochemistry: Electronic Structure Methods and Force Analysis. *Chem. Rev.* **2016**, *116*, 14137–14180. [[CrossRef](#)]
13. McNaught, A.D.; Wilkinson, A. *IUPAC. Compendium of Chemical Technology*, 2nd ed.; Blackwell Scientific Publications: Oxford, UK, 1997; ISBN 0865426848.
14. Urakaev, F.K.; Boldyrev, V.V. Mechanism and kinetics of mechanochemical processes in comminuting devices: 1. Theory. *Powder Technol.* **2000**, *107*, 93–107. [[CrossRef](#)]
15. Urakaev, F.K.; Boldyrev, V.V. Mechanism and kinetics of mechanochemical processes in comminuting devices: 2. Applications of the theory. Experiment. *Powder Technol.* **2000**, *107*, 197–206. [[CrossRef](#)]
16. Boldyrev, V.V.; Tkáčová, K. Mechanochemistry of solids: Past, Present and Prospects. *J. Mater. Synth. Process.* **2000**, *8*, 121–132. [[CrossRef](#)]
17. Cintas, P.; Tagliapietra, S.; Caporaso, M.; Tabasso, S.; Cravotto, G. Enabling Technologies Built on a Sonochemical Platform: Challenges and Opportunities. *Ultrason. Sonochem.* **2015**, *25*, 8–16. [[CrossRef](#)]
18. Baláž, P.; Achimovičová, M.; Baláž, M.; Billik, P.; Cherkezova-Zheleva, Z.; Criado, J.M.; Delogu, F.; Dutková, E.; Gaffet, E.; Gotor, F.J.; et al. Hallmarks of Mechanochemistry: From Nanoparticles to Technology. *Chem. Soc. Rev.* **2013**, *42*, 7571–7637. [[CrossRef](#)]
19. Dolotko, O.; Wiench, J.W.; Dennis, K.W.; Pecharsky, V.K.; Balema, V.P. Mechanically induced reactions in organic solids: Liquid eutectics or solid-state processes? *New J. Chem.* **2010**, *34*, 25–28. [[CrossRef](#)]
20. Zhang, H.; Lin, Y.; Xu, Y.; Weng, W. Mechanochemistry of Topological Complex Polymer Systems. *Top. Curr. Chem.* **2015**, *369*, 135–207. [[CrossRef](#)]
21. Cintas, P.; Cravotto, G.; Barge, A.; Martina, K. Interplay between Mechanochemistry and Sonochemistry. In *Polymer Mechanochemistry*; Boulatov, R., Ed.; Springer International Publishing: Cham, Switzerland, 2015; pp. 239–284. ISBN 978-3-319-22825-9.
22. Michalchuk, A.A.L.; Tumanov, I.A.; Boldyreva, E.V. Complexities of mechanochemistry: Elucidation of processes occurring in mechanical activators via implementation of a simple organic system. *CrystEngComm* **2013**, *15*, 6403–6412. [[CrossRef](#)]
23. Wang, J. Characterization and Applications of Force-Induced Reactions. Ph.D. Thesis, Duke University, Durham, NC, USA, 2015.
24. Haruta, N.; Sato, T.; Tanaka, K.; Baron, M. Reaction mechanism in the mechanochemical synthesis of dibenzophenazine: Application of vibronic coupling density analysis. *Tetrahedron Lett.* **2013**, *54*, 5920–5923. [[CrossRef](#)]
25. Oliveira, P.F.M.; Baron, M.; Chamayou, A.; Baltas, M.; Guidetti, B.; Haruta, N.; Tanaka, K.; Sato, T. Lowering the Activation Energy under Mechanochemical Conditions: The Case of 2,3-diphenylquinoxaline. *ChemistrySelect* **2016**, *1*, 984–988. [[CrossRef](#)]
26. Ribas-Arino, J.; Shiga, M.; Marx, D. Understanding Covalent Mechanochemistry. *Angew. Chem. Int. Ed.* **2009**, *48*, 4190–4193. [[CrossRef](#)]
27. Haruta, N.; Oliveira, P.F.M.; Sato, T.; Tanaka, K.; Baron, M. Force-Induced Dissolution of Imaginary Mode in Mechanochemical Reaction: Dibenzophenazine Synthesis. *J. Phys. Chem. C* **2019**, *123*, 21581–21587. [[CrossRef](#)]
28. Smalø, H.S.; Rybkin, V.V.; Klopper, W.; Helgaker, T.; Uggerud, E. Mechanochemistry: The Effect of Dynamics. *J. Phys. Chem. A* **2014**, *118*, 7683–7694. [[CrossRef](#)]
29. Friščić, T.; Jones, W. Recent Advances in Understanding the Mechanism of Cocrystal Formation via Grinding. *Cryst. Growth Des.* **2009**, *9*, 1621–1637. [[CrossRef](#)]

30. Bonnamour, J.; Métro, T.X.; Martinez, J.; Lamaty, F. Environmentally Benign Peptide Synthesis Using Liquid-Assisted Ball-Milling: Application to the Synthesis of Leu-Enkephalin. *Green Chem.* **2013**, *15*, 1116–1120. [[CrossRef](#)]
31. Tumanov, I.A.; Achkasov, A.F.; Boldyreva, E.V.; Boldyrev, V.V. Following the products of mechanochemical synthesis step by step. *CrystEngComm* **2011**, *13*, 2213–2216. [[CrossRef](#)]
32. Užarević, K.; Štrukil, V.; Mottillo, C.; Julien, P.A.; Puškarić, A.; Friščić, T.; Halasz, I. Exploring the Effect of Temperature on a Mechanochemical Reaction by in Situ Synchrotron Powder X-ray Diffraction. *Cryst. Growth Des.* **2016**, *16*, 2342–2347. [[CrossRef](#)]
33. Kubota, K.; Takahashi, R.; Ito, H. Mechanochemistry Allows Carrying out Sensitive Organometallic Reactions in Air: Glove-Box-and-Schlenk-Line-Free Synthesis of Oxidative Addition Complexes from Aryl Halides and Palladium(0). *Chem. Sci.* **2019**, *10*, 5837–5842. [[CrossRef](#)] [[PubMed](#)]
34. Ingner, F.J.L.; Giustra, Z.X.; Novosedlik, S.; Orthaber, A.; Gates, P.J.; Dyrager, C.; Pilarski, L.T. Mechanochemical Synthesis of (Hetero)Aryl Au(I) Complexes. *Green Chem.* **2020**, *22*, 5648–5655. [[CrossRef](#)]
35. Takacs, L. The Historical Development of Mechanochemistry. *Chem. Soc. Rev.* **2013**, *42*, 7649–7659. [[CrossRef](#)]
36. Ribas-Arino, J.; Marx, D. Covalent Mechanochemistry: Theoretical Concepts and Computational Tools with Applications to Molecular Nanomechanics. *Chem. Rev.* **2012**, *112*, 5412–5487. [[CrossRef](#)]
37. Toda, F.; Tanaka, K.; Iwata, S. Oxidative Coupling Reactions of Phenols with Iron(III) Chloride in the Solid State. *J. Org. Chem.* **1989**, *54*, 3007–3009. [[CrossRef](#)]
38. Kaupp, G. Solvent-Less Organic Synthesis. In *Kirk-Othmer Encyclopedia of Chemical Technology*; Othmer, K., Ed.; Wiley Online Library: Cambridge, MA, USA, 2012. [[CrossRef](#)]
39. Baron, M. Towards a Greener Pharmacy by More Eco Design. *Waste Biomass Valorization* **2012**, *3*, 395–407. [[CrossRef](#)]
40. Achar, T.K.; Bose, A.; Mal, P. Mechanochemical Synthesis of Small Organic Molecules. *Beilstein, J. Org. Chem.* **2017**, *13*, 1907–1931. [[CrossRef](#)] [[PubMed](#)]
41. Tanaka, K.; Toda, F. Solvent-Free Organic Synthesis. *Chem. Rev.* **2000**, *100*, 1025–1074. [[CrossRef](#)]
42. Carlier, L.; Baron, M.; Chamayou, A.; Couarraze, G. Use of co-grinding as a solvent-free solid state method to synthesize dibenzophenazines. *Tetrahedron Lett.* **2011**, *52*, 4686–4689. [[CrossRef](#)]
43. Carlier, L.; Baron, M.; Chamayou, A.; Couarraze, G. Greener pharmacy using solvent-free synthesis: Investigation of the mechanism in the case of dibenzophenazine. *Powder Technol.* **2013**, *240*, 41–47. [[CrossRef](#)]
44. Konnert, L.; Gauliard, A.; Lamaty, F.; Martinez, J.; Colacino, E. Solventless Synthesis of N-Protected Amino Acids in a Ball Mill. *ACS Sustain. Chem. Eng.* **2013**, *1*, 1186–1191. [[CrossRef](#)]
45. Mashkouri, S.; Naimi-Jamal, M.R. Mechanochemical solvent-free and catalyst-free one-pot synthesis of pyrano[2,3-d]pyrimidine-2,4(1H,3H)-diones with quantitative yields. *Molecules* **2009**, *14*, 474–479. [[CrossRef](#)]
46. Métro, T.X.; Bonnamour, J.; Reidon, T.; Sarpoulet, J.; Martinez, J.; Lamaty, F. Mechanochemical synthesis of amides in the total absence of organic solvent from reaction to product recovery. *Chem. Commun.* **2012**, *48*, 11781–11783. [[CrossRef](#)]
47. Ranu, B.; Stolle, A. *Ball Milling Towards Green Synthesis: Applications, Projects, Challenges*; Royal Society of Chemistry: Cambridge, UK, 2015; ISBN 978-1-84973-945-0.
48. Rodríguez, B.; Rantanen, T.; Bolm, C. Solvent-free asymmetric organocatalysis in a ball mill. *Angew. Chem. Int. Ed.* **2006**, *118*, 7078–7080. [[CrossRef](#)]
49. Stolle, A.; Szuppa, T.; Leonhardt, S.E.S.; Ondruschka, B. Ball milling in organic synthesis: Solutions and challenges. *Chem. Soc. Rev.* **2011**, *40*, 2317–2329. [[CrossRef](#)]
50. Štrukil, V.; Bartolec, B.; Portada, T.; Đilović, I.; Halasz, I.; Margetić, D. One-pot mechanochemical synthesis of aromatic amides and dipeptides from carboxylic acids and amines. *Chem. Commun.* **2012**, *48*, 12100–12102. [[CrossRef](#)]
51. Zhu, X.; Li, Z.; Jin, C.; Xu, L.; Wu, Q.; Su, W. Mechanically activated synthesis of 1,3,5-triaryl-2-pyrazolines by high speed ball milling. *Green Chem.* **2009**, *11*, 163–165. [[CrossRef](#)]
52. Qiu, Y.; Puni, K.T.; Duplan, C.C.; Lindsay, A.C.; Sperry, J. Tetrahydrocarbazoles by mechanochemical Fisher indolization. *Tetrahedron Lett.* **2021**, *72*, 153068. [[CrossRef](#)]
53. Nibin Joy, M.; Bodke, Y.D.; Telkar, S.; Bakulev, V.A. Synthesis of Coumarins Linked With 1,2,3-Triazoles under Microwave Irradiation and Evaluation of their Antimicrobial and Antioxidant Activity. *J. Mex. Chem. Soc.* **2020**, *64*, 53–73. [[CrossRef](#)]
54. Nibin Joy, M.; Beliaev, N.; Beryozkina, T.V.; Bakulev, V.A. Design and the synthesis of 1-heteroaryl-1,2,3-triazoles connected to coumarins via ether linker. *J. Heterocycl. Chem.* **2020**, *57*, 3173–3185. [[CrossRef](#)]
55. Rishikesan, R.; Ranjith, P.K.; Nibin Joy, M.; Ayyiliath, M.S.; Koti Reddy, E.; Rajeesh, P.; Karickal, R.H.; Vaishnav, B.; Kereyagalally, H.N.; Muralidharan, A. Synthesis of some novel piperidine fused 5-thioxo-1H-1,2,4-triazoles as potential antimicrobial and antitubercular agents. *J. Chem. Sci.* **2021**, *133*, 3. [[CrossRef](#)]
56. Dheer, D.; Singh, V.; Shankar, R. Medicinal Attributes of 1,2,3-Triazoles: Current Developments. *Bioorganic Chem.* **2017**, *71*, 30–54. [[CrossRef](#)] [[PubMed](#)]
57. Alexandre, J.A.C.; Swan, M.K.; Latchem, M.J.; Boyall, D.; Pollard, J.R.; Hughes, S.W.; Westcott, J. New 4-Amino-1,2,3-Triazole Inhibitors of Indoleamine 2,3-Dioxygenase Form a Long-Lived Complex with the Enzyme and Display Exquisite Cellular Potency. *ChemBioChem* **2018**, *19*, 552–561. [[CrossRef](#)] [[PubMed](#)]
58. Wu, G.; Gao, Y.; Kang, D.; Huang, B.; Huo, Z.; Liu, H.; Poongavanam, V.; Zhan, P.; Liu, X. Design, Synthesis and Biological Evaluation of Tacrine-1,2,3-Triazole Derivatives as Potent Cholinesterase Inhibitors. *MedChemComm* **2018**, *9*, 149–159. [[CrossRef](#)] [[PubMed](#)]

59. Ashok, D.; Chiranjeevi, P.; Kumar, A.V.; Sarasija, M.; Krishna, V.S.; Sriram, D.; Balasubramanian, S. 1,2,3-Triazole-Fused Spirochromenes as Potential Anti-Tubercular Agents: Synthesis and Biological Evaluation. *RSC Adv.* **2018**, *8*, 16997–17007. [[CrossRef](#)]
60. López-Rojas, P.; Janeczko, M.; Kubiński, K.; Amesty, Á.; Masłyk, M.; Estévez-Braun, A.; López-Rojas, P.; Janeczko, M.; Kubiński, K.; Amesty, Á.; et al. Synthesis and Antimicrobial Activity of 4-Substituted 1,2,3-Triazole-Coumarin Derivatives. *Molecules* **2018**, *23*, 199. [[CrossRef](#)]
61. Mady, M.F.; Awad, G.E.A.; Jørgensen, K.B. Ultrasound-Assisted Synthesis of Novel 1,2,3-Triazoles Coupled Diaryl Sulfone Moieties by the CuAAC Reaction, and Biological Evaluation of Them as Antioxidant and Antimicrobial Agents. *Eur. J. Med. Chem.* **2014**, *84*, 433–443. [[CrossRef](#)]
62. Nallapati, S.B.; Sreenivas, B.Y.; Bankala, R.; Parsa, K.V.L.; Sripelly, S.; Mukkanti, K.; Pal, M. 1,2,3-Triazoles Derived from Olanzapine: Their Synthesis via an Ultrasound Assisted CuAAC Method and Evaluation as Inhibitors of PDE4B. *RSC Adv.* **2015**, *5*, 94623–94628. [[CrossRef](#)]
63. Rezki, N. A Green Ultrasound Synthesis, Characterization and Antibacterial Evaluation of 1,4-Disubstituted 1,2,3-Triazoles Tethering Bioactive Benzothiazole Nucleus. *Molecules* **2016**, *21*, 505. [[CrossRef](#)]
64. Rezki, N.; Aouad, M.R. Green Ultrasound-Assisted Three-Component Click Synthesis of Novel 1H-1,2,3-Triazole Carrying Benzothiazoles and Fluorinated-1,2,4-Triazole Conjugates and Their Antimicrobial Evaluation. *Acta Pharm. Zagreb Croat.* **2017**, *67*, 309–324. [[CrossRef](#)]
65. Vadivelu, M.; Sugirdha, S.; Dheenkumar, P.; Arun, Y.; Karthikeyan, K.; Praveen, C. Solvent-Free Implementation of Two Dissimilar Reactions Using Recyclable CuO Nanoparticles under Ball-Milling Conditions: Synthesis of New Oxindole-Triazole Pharmacophores. *Green Chem.* **2017**, *19*, 3601–3610. [[CrossRef](#)]
66. Mack, J.; Shumba, M. Rate Enhancement of the Morita–Baylis–Hillman Reaction through Mechanochemistry. *Green Chem.* **2007**, *9*, 328–330. [[CrossRef](#)]
67. Rubio, N.; Mei, K.C.; Klippstein, R.; Costa, P.M.; Hodgins, N.; Wang, J.T.W.; Festy, F.; Abbate, V.; Hider, R.C.; Chan, K.L.A.; et al. Solvent-Free Click-Mechanochemistry for the Preparation of Cancer Cell Targeting Graphene Oxide. *ACS Appl. Mater. Interfaces* **2015**, *7*, 18920–18923. [[CrossRef](#)] [[PubMed](#)]
68. Liu, L.; Zhang, S.; Xue, F.; Lou, G.; Zhang, H.; Ma, S.; Duan, W.; Wang, W. Catalytic Enantioselective Henry Reactions of Isatins: Application in the Concise Synthesis of (S)-(–)-Spirobrassinin. *Chem.—Eur. J.* **2011**, *17*, 7791–7795. [[CrossRef](#)] [[PubMed](#)]
69. DeLorbe, J.E.; Horne, D.; Jove, R.; Mennen, S.M.; Nam, S.; Zhang, F.L.; Overman, L.E. General Approach for Preparing Epidithiodioxopiperazines from Trioxopiperazine Precursors: Enantioselective Total Syntheses of (+)- and (–)-Gliocladine C, (+)-Leptosin D, (+)-T988C, (+)-Bioneectin A, and (+)-Gliocladin A. *J. Am. Chem. Soc.* **2013**, *135*, 4117–4128. [[CrossRef](#)] [[PubMed](#)]
70. Mohammadi, S.; Heiran, R.; Herrera, R.P.; Marqués-López, E. Isatin as a Strategic Motif for Asymmetric Catalysis. *ChemCatChem* **2013**, *5*, 2131–2148. [[CrossRef](#)]
71. Sahu, A.; Agrawal, R.K.; Pandey, R. Synthesis and Systemic Toxicity Assessment of Quinine-Triazole Scaffold with Antiprotozoal Potency. *Bioorganic Chem.* **2019**, *88*, 102939. [[CrossRef](#)]
72. Blasco, B.; Leroy, D.; Fidock, D.A. Antimalarial Drug Resistance: Linking Plasmodium Falciparum Parasite Biology to the Clinic. *Nat. Med.* **2017**, *23*, 917–928. [[CrossRef](#)]
73. Adarsh, S.; Deepak, K.; Ram, K.A. Antileishmanial Drug Discovery: Synthetic Methods, Chemical Characteristics, and Biological Potential of Quinazolines and Its Derivatives. *Anti-Inflamm. Anti-Allergy Agents Med. Chem.* **2017**, *16*, 3–32.
74. Scriven, E.F.V.; Turnbull, K. Azides: Their Preparation and Synthetic Uses. *Chem. Rev.* **1988**, *88*, 297–368. [[CrossRef](#)]
75. Sampath, S.; Vadivelu, M.; Ravindran, R.; Perumal, P.T.; Velkannan, V.; Karthikeyan, K. Synthesis of 1,2,3-Triazole Tethered 3-Hydroxy-2-Oxindoles: Promising Corrosion Inhibitors for Steel in Acidic Medium and Their Anti-Microbial Evaluation. *ChemistrySelect* **2020**, *5*, 2130–2134. [[CrossRef](#)]
76. Singh, G.S.; Desta, Z.Y. Isatins as Privileged Molecules in Design and Synthesis of Spiro-Fused Cyclic Frameworks. *Chem. Rev.* **2012**, *112*, 6104–6155. [[CrossRef](#)]
77. Hong, L.; Wang, R. Recent Advances in Asymmetric Organocatalytic Construction of 3,3'-Spirocyclic Oxindoles. *Adv. Synth. Catal.* **2013**, *355*, 1023–1052. [[CrossRef](#)]
78. Trost, B.M.; Brennan, M.K. Asymmetric Syntheses of Oxindole and Indole Spirocyclic Alkaloid Natural Products. *Synthesis* **2009**, *2009*, 3003–3025. [[CrossRef](#)]
79. Williams, R.M.; Cox, R.J. Paraherquamides, Brevianamides, and Asperparalines: Laboratory Synthesis and Biosynthesis. An Interim Report. *Acc. Chem. Res.* **2003**, *36*, 127–139. [[CrossRef](#)]
80. Satyamaheshwar, P. 3-Substituted-3-Hydroxy-2-Oxindole, an Emerging New Scaffold for Drug Discovery with Potential Anti-Cancer and Other Biological Activities. *Curr. Bioact. Compd.* **2009**, *5*, 20–38. [[CrossRef](#)]
81. Rottmann, M.; McNamara, C.; Yeung, B.K.S.; Lee, M.C.S.; Zou, B.; Russell, B.; Seitz, P.; Plouffe, D.M.; Dharia, N.V.; Tan, J.; et al. Spiroindolones, a Potent Compound Class for the Treatment of Malaria. *Science* **2010**, *329*, 1175–1180. [[CrossRef](#)]
82. Khetmalis, Y.M.; Shivani, M.; Murugesan, S.; Chandra Sekhar, K.V.G. Oxindole and Its Derivatives: A Review on Recent Progress in Biological Activities. *Biomed. Pharmacother.* **2021**, *141*, 111842. [[CrossRef](#)] [[PubMed](#)]
83. Clubley, B.G. *Chemical Inhibitors for Corrosion*; Control, Royal Society Of Chemistry: Cambridge, UK, 1990; ISBN 978-0851868363.
84. Sastri, V.S. *Corrosion Inhibitors: Principles and Applications*; Wiley: New York, NY, USA, 1998; ISBN 9780471976080.

85. Zhou, C.H.; Wang, Y. Recent Researches in Triazole Compounds as Medicinal Drugs. *Curr. Med. Chem.* **2012**, *19*, 239–280. [[CrossRef](#)] [[PubMed](#)]
86. Tennefy, A.B. *Pesticide Research Trends*, 14th ed.; Nova Science Publishers: New York, NY, USA, 2008; ISBN 978-1604562002.
87. Nakka, M.; Tadikonda, R.; Rayavarapu, S.; Sarakula, P.; Vidavalur, S. A Simple and Efficient Synthesis of 3,4,5-Trisubstituted/N-Fused 1,2,4-Triazoles via Ceric Ammonium Nitrate Catalyzed Oxidative Cyclization of Amidrazones with Aldehydes Using Polyethylene Glycol as a Recyclable Reaction Medium. *Synthesis* **2015**, *47*, 517–525. [[CrossRef](#)]
88. Szócs, B.; Bokor, É.; Szabó, K.E.; Kiss-Szikszai, A.; Tóth, M.; Somsák, L. Synthesis of 5-Aryl-3-C-Glycosyl- and Unsymmetrical 3,5-Diaryl-1,2,4-Triazoles from Alkylidene-Amidrazones. *RSC Adv.* **2015**, *5*, 43620–43629. [[CrossRef](#)]
89. Aly, A.A.; Hassan, A.A.; Mohamed, N.K.; Ramadadfon, M.; Brown, A.B.; Abd El-Aal, A.S.; Bräse, S.; Nieger, M. Regioselective Formation of 1,2,4-Triazoles by the Reaction of Amidrazones in the Presence of Diethyl Azodicarboxylate and Catalyzed by Triethylamine. *Mol. Divers.* **2019**, *23*, 195–203. [[CrossRef](#)] [[PubMed](#)]
90. Singh, R.; Kashaw, S.K.; Mishra, V.K.; Mishra, M.; Rajoriya, V.; Kashaw, V. Design and Synthesis of New Bioactive 1,2,4-Triazoles, Potential Antitubercular and Antimicrobial Agents. *Indian J. Pharm. Sci.* **2018**, *80*, 36–45. [[CrossRef](#)]
91. Sonawane, A.D.; Rode, N.D.; Nawale, L.; Joshi, R.R.; Joshi, R.A.; Likhite, A.P.; Sarkar, D. Synthesis and Biological Evaluation of 1,2,4-Triazole-3-Thione and 1,3,4-Oxadiazole-2-Thione as Antimycobacterial Agents. *Chem. Biol. Drug Des.* **2017**, *90*, 200–209. [[CrossRef](#)] [[PubMed](#)]
92. Liu, C.F.; Shen, Q.K.; Li, J.J.; Tian, Y.S.; Quan, Z. Synthesis and Biological Evaluation of Novel 7-Hydroxy-4-Phenylchromen-2-One-Linked to Triazole Moieties as Potent Cytotoxic Agents. *J. Enzyme Inhib. Med. Chem.* **2017**, *32*, 1111–1119. [[CrossRef](#)]
93. De, P.; Koumba Yoya, G.; Constant, P.; Bedos-Belval, F.; Duran, H.; Saffon, N.; Daffé, M.; Baltas, M. Design, Synthesis, and Biological Evaluation of New Cinnamic Derivatives as Antituberculosis Agents. *J. Med. Chem.* **2011**, *54*, 1449–1461. [[Cross-Ref](#)]
94. Veau, D.; Krykun, S.; Mori, G.; Orena, B.S.; Pasca, M.R.; Frongia, C.; Lobjois, V.; Chassaing, S.; Lherbet, C.; Baltas, M. Triazolophthalazines: Easily Accessible Compounds with Potent Antitubercular Activity. *ChemMedChem* **2016**, *11*, 1078–1089. [[CrossRef](#)]
95. Gonnet, L.; Guidetti, B.; André-Barrès, C.; Calvet, R.; Maturano, M.; Menendez, C.; Lherbet, C.; Baron, M.; Chamayou, A.; Baltas, M. Study of the Two Steps and One-Pot Two-Step Mechanochemical Synthesis of Annulated 1,2,4-Triazoles. *ACS Sustain. Chem. Eng.* **2020**, *8*, 3114–3125. [[CrossRef](#)]
96. Gonnet, L. Mécanosynthèses Organiques: Étude Cinétique de la Réaction de Diels-Alder et Synthèses de 1,2,4-Triazoles à Activités Biologiques. Ph.D. Thesis, Ecole des Mines d'Albi-Carmaux, Albi, France, 2019. (In French).
97. Wang, Q.; Chittaboina, S.; Barnhill, H.N. Advances in 1,3-Dipolar Cycloaddition Reaction of Azides and Alkynes—A Prototype of “Click” Chemistry. *Lett. Org. Chem.* **2005**, *2*, 293–301. [[CrossRef](#)]
98. Brantley, J.N.; Wiggins, K.M.; Bielawski, C.W. Unclicking the Click: Mechanically Facilitated 1,3-Dipolar Cycloreversions. *Science* **2011**, *333*, 1606–1609. [[CrossRef](#)] [[PubMed](#)]
99. McNutt, M. Editorial Retraction. *Science* **2015**, *347*, 834. [[CrossRef](#)] [[PubMed](#)]
100. Brantley, J.N. Mechanical Activations of Synthetic and Biological Systems. Ph.D. Thesis, University of Texas, Austin, TX, USA, 2014.
101. Brantley, J.N.; Konda, S.S.M.; Makarov, D.E.; Bielawski, C.W. Regiochemical Effects on Molecular Stability: A Mechanochemical Evaluation of 1,4- and 1,5-Disubstituted Triazoles. *J. Am. Chem. Soc.* **2012**, *134*, 9882–9885. [[CrossRef](#)] [[PubMed](#)]
102. Stauch, T.; Dreuw, A. Force-Induced Retro-Click Reaction of Triazoles Competes with Adjacent Single-Bond Rupture. *Chem. Sci.* **2017**, *8*, 5567–5575. [[CrossRef](#)] [[PubMed](#)]
103. Krupička, M.; Dopieralski, P.; Marx, D. Unclicking the Click: Metal-Assisted Mechanochemical Cycloreversion of Triazoles Is Possible. *Angew. Chem. Int. Ed.* **2017**, *56*, 7745–7749. [[CrossRef](#)] [[PubMed](#)]